ESMO Congress, Barcelona, September 27 – October 1, 2019 Abstract 3340

Corresponding author e-mail: jealesm@gmail.com

Effectiveness of olaparib plus trastuzumab in patients with HER2-positive, germinal *BRCA*–mutated or Homologous Recombination Deficient advanced breast cancer. The OPHELIA Study.

J.E. Alés-Martínez[1], S. Morales[2], M. Fernández Abad[3], P. Sánchez-Rovira[4], F.J. Salvador Bofill[5], A. Lahuerta[6], J.A. García-Sáenz[7], I. Garau Llinas[8], T. Díaz Redondo[9], N. Ferrer[10], V. Carañana[11], R. López[12], S.E. Alonso Soler[13], B. Bermejo[14], J. de la Haba[15], P. Zamora[16], J. Balmaña[17], E. López-Miranda[18], J. Cortés[19], A. Llombart Cussac[20];

[1] Medical Oncology, Hospital Nuestra Señora de Sonsoles, Ávila, Spain [2]Medical Oncology, Hospital Universitari Arnau de Vilanova de Lleida, Lleida, Spain, [3]Medical Oncology, Hospital Universitario Ramón y Cajal, Madrid, Spain [4]Medical Oncology, Complejo Hospitalario de Jaén, Jaén, Spain, [5]Medical Oncology, Hospital Universitario Virgen del Rocío, Sevilla, Spain, [6]Medical Oncology, Fundación Onkologikoa, Donostia-San Sebastián, Spain [7]Medical Oncology, Hospital Clínico San Carlos, Madrid, Spain [8]Medical Oncology, Hospital Son Llatzer, Palma De Mallorca, Spain [9]Medical Oncology, Hospital Universitario Virgen de la Victoria, Malaga, Spain [10]Medical Oncology, Hospital Universitari Son Espases, Palma De Mallorca, Spain,[11]Medical Oncology, Hospital Arnau de Vilanova, Valencia, Spain, [12]Medical Oncology, Hospital Clínico Universitario de Santiago (CHUS), Santiago De Compostela, Spain [13]Medical Oncology, Hospital San Pedro de Alcántara, Cáceres, Spain [14]Medical Oncology, Hospital Clínic Universitari de València, Valencia, Spain, [15]Medical Oncology, Hospital Universitario Reina Sofía, Córdoba, Spain, [16]Medical Oncology, Hospital Universitario La Paz, Madrid, Spain [17]Medical Oncology, Vall d'Hebron University Hospital, Barcelona, Spain, [18]Medical Oncology, Medica Scientia Innovation Research (MedSIR), Ridgewood, New Jersey, USA & Barcelona, Spain; Hospital Universitario Ramón y Cajal, Madrid, Spain [19]Medical Oncology, Vall d´Hebron Institute of Oncology (VHIO), Barcelona, Spain; IOB Institute of Oncology, Quironsalud Group, Madrid and Barcelona, Spain; Medica Scientia Innovation Research (MedSIR), Ridgewood, New Jersey, USA, & Barcelona, Spain, [20]Medical Oncology, Medica Scientia Innovation Research (MedSIR), Ridgewood, New Jersey, USA & Barcelona, Spain; FISABIO, Hospital Arnau de Vilanova, Valencia, Spain

BACKGROUND

- In the OlympiAD trial the use of poly(adenosine diphosphate–ribose) polymerase inhibitor (PARPi) olaparib (Lynparza®) resulted in better progression–free survival (PFS) than standard hemotherapy, which led to approval in patients (pts) with Human Epidermal Growth Factor Receptor 2 (HER2)-negative, germline *BRCA1*- and/or *BRCA2*-mutated (gBRCAm) metastatic breast cancer [(1)].
- There is a potential therapeutic role for PARPis in pts with defective DNA repair [(2)].
- Preclinical data support a strong synergism between olaparib (Lynparza®) and trastuzumab in HER2-positive (HER2[+]) cells that are sensitive to PARPis [(3)].
- This study will evaluate the efficacy and safety of olaparib (Lynparza®) plus trastuzumab in pts with HER2[+], gBRCAm (Cohort A) or germline *BRCA1*- and/or *BRCA2*-wild-type (gBRCAwt) / Homologous Recombination Deficient (HRD) (Cohort B) advanced breast cancer (ABC).

OBJECTIVES

- **PRIMARY OBJECTIVE**

To assess the efficacy –as determined by the overall response rate (ORR) and progression-free survival (PFS) based on Response evaluation criteria in solid tumors (RECIST) v.1.1– of olaparib (Lynparza®) in combination with trastuzumab in pts with HER2[+], gBRCAm ABC (Cohort A).

- **CO-PRIMARY ENDPOINTS**
  - ORR is defined as the percentage of pts with complete response (CR) or partial response (PR) confirmed at least 4 weeks after the initial response assessment.
  - PFS is defined as the time from treatment initiation to the date of the first documented progressive disease or death due to any cause, whichever occurs first.

- **SECONDARY OBJECTIVES**
  - To assess the efficacy –as determined by the clinical benefit rate (CBR), duration of response (DoR), maximum tumor reduction and overall survival (OS)– of olaparib (Lynparza®) in combination with trastuzumab in pts with HER2[+], gBRCAm ABC (Cohort A).
  - To assess the efficacy –as determined by the ORR, PFS, CBR, DoR, maximum tumor reduction and OS– of olaparib (Lynparza®) in combination with trastuzumab in pts with HER2[+], gBRCAwt / HRD ABC (Cohort B)
  - To evaluate health-related quality of life (HRQoL) measured by the European Organization for Research and Treatment of Cancer (EORTC) and EuroQoL 5D (EQ-5D) questionnaires both in Cohort A and Cohort B.

- **SAFETY OBJECTIVES**
  - To assess adverse events assessed using the National Cancer Institute's Common Terminology Criteria for Adverse Events (NCI CTCAE) v5.0.

- **EXPLORATORY OBJECTIVES**
  - To assess prevalence and predictive value of both genomic profiles (mutations in germline *BRCA1* and/or *BRCA2* genes and gBRCAwt / HRD status) in HER2[+] ABC pts.
  - To investigate new molecular markers that are predictive of response and/or resistance to olaparib (Lynparza®) in combination with trastuzumab.

TRIAL DESIGN

- This is a multicenter, single–arm, two–cohort, Simon's two–stage, phase II trial.
- The cohort A will recruit N=20 gBRCAm ABC pts. The cohort B will recruit N=13 gBRCAwt / HRD ABC pts that will require prospective central testing of both tumor tissue and blood samples according to the whole genome sequencing (WGS)-based HRDetect predictor at the Wellcome Trust Sanger Institute [(4)].
- Pts will receive olaparib (Lynparza®) tablets (300mg taken oral twice daily during 21–day cycles) in combination with trastuzumab (intravenous or subcutaneous at standard dose) until progression or unacceptable toxicity.

OPHELIA Study Design



ABC: Advance breast cancer; AEs: Adverse events; BID: Twice daily administration; CBR: Clinical benefit rate; CHT: Chemotherapy; CR: Complete response; DoR: Duration of response; ECOG: Eastern Cooperative Oncology Group; FISH: Fluorescence in situ hybridization; gBRCAm: germline BRCA1- and/or BRCA2-mutated; HER2[+]: Human Epidermal Growth Factor Receptor 2-positive; HRD: Homologous Recombination Deficient; HRQoL: Health-related quality of life; IHC: Immunohistochemistry; mo.: Months; IV: Intravenous; ORR: Overall response rate; PARPi: Poly(adenosine diphosphate–ribose) polymerase inhibitor; PD: Progressive disease; PFS: Progression-free survival; PO: Per os, orally; PR: Partial response; pts: Patients; RECIST: Response evaluation criteria in solid tumors; SC: Subcutaneous.

* gBRCA / HRD status (negrita): Prospective central testing of both tumor tissue and blood samples according to the whole genome sequencing-based HRDetect predictor at the Wellcome Trust Sanger Institute [(4)].

- **SAMPLE SIZE**

· Cohort A

We plan a Simon's minimax two–stage design, with 7 pts in stage I and, if ≥1 responding pts, additional 13 pts in stage II.

1st co-primary endpoint (ORR):

Final ORR will be promising with ≥4 responders among 18 evaluable pts (H0: ORR≤5%; HA: ORR≥30%). These give us an 80% power at 0.025 one–sided alpha level. We will accrual 20 pts to account a 10% drop-out rate correction.

Second co-primary endpoint (PFS):

PFS estimation in Cohort A will be based a one–arm log–rank test (H0: PFS≤3–months; HA: PFS≥6–months). These give us an 80% power at 0.025 one–sided alpha level.

· Exploratory Cohort B

Additional 13 pts in stage II will be recruited according to a binomial one-stage design.

Final ORR will be promising with ≥3 responders among 13 pts (p<0.025; H0: ORR≤5%).

BIBLIOGRAPHY

(1) Robson M, et al. *N Engl J Med*. 2017 Aug 10;377(6):523-533.
(2) Mirza MR et al. *N Engl J Med*. 2016 Dec 1;375(22):2154–64
(3) García-Parra J et al. *Eur J Cancer*. 2014 Oct;50(15):2725–34.
(4) Davies H. et al. *Nat Med*. 2017 Apr;23(4):517-525.

Conflicts of interest: https://medsir.org/ophelia

ACKNOWLEDGEMENTS:
We thank all participating patients and study team involved in **OPHELIA trial**. This study was funded by AstraZeneca.

Trial registration: NCT03931551. **Date of registration:** April 30, 2019. **First patient included:** August 7, 2019.

Copies of this poster obtained through QR and/or text key codes are for personal use only and may not be reproduced without written permission of the authors.